arXiv:nlin/0412050v2 [nlin.SI] 3 Mar 2005

Soliton Solutions for the Non-autonomous Discrete-time Toda Lattice Equation

Kenji Kajiwara[1,2] and Atsushi Mukaihira[1]

[1]: Graduate School of Mathematics, Kyushu University,
6-10-1 Hakozaki, Fukuoka 812-8581, Japan

[2]: School of Mathematics and Statistics F07, University of Sydney,
Sydney, NSW 2006, Australia

Abstract

We construct N-soliton solution for the non-autonomous discrete-time Toda lattice equation, which is a generalization of the discrete-time Toda equation such that the lattice interval with respect to time is an arbitrary function in time.

1 Introduction

In this article, we consider the nonlinear partial difference equation given by

$$\begin{aligned} &A_n^{t+1} + B_n^{t+1} + \lambda_{t+1} = A_n^t + B_{n+1}^t + \lambda_t, \\ &A_{n-1}^{t+1} B_n^{t+1} = A_n^t B_n^t, \end{aligned} \qquad n \in \mathbb{Z}, \tag{1}$$

where n and t are the independent variables, A_n^t and B_n^t are the dependent variables and λ_t is an arbitrary function in t, respectively. In the physical context, the variables n, t, A_n^t and B_n^t are identical as the lattice site, the discrete time and the fields, respectively. Equation (1) is equivalent to the following equation

$$\begin{aligned} &J_n^{t+1} - \delta_{t+1} V_{n-1}^{t+1} = J_n^t - \delta_t V_n^t, \\ &V_n^{t+1}(1 - \delta_{t+1} J_n^{t+1}) = V_n^t (1 - \delta_t J_{n+1}^t), \end{aligned} \qquad n \in \mathbb{Z}, \tag{2}$$

where the variables are related as

$$A_n^t = -\lambda_t + J_n^t, \quad B_n^t = -\lambda_t^{-1} V_{n-1}^t, \quad \delta_t = \lambda_t^{-1}, \tag{3}$$

respectively. When λ_t or δ_t is a constant, equation (1) or (2) reduces to the discrete-time Toda equation proposed by Hirota [2, 4]. Moreover, equation (2) yields the celebrated Toda lattice equation

$$\begin{aligned} &\frac{dJ_n}{dt} = V_{n-1} - V_n, \\ &\frac{dV_n}{dt} = V_n (J_n - J_{n+1}), \end{aligned} \qquad n \in \mathbb{Z}, \tag{4}$$

in the continous limit $\delta_t = \delta \to 0$.

Equation (1) was proposed by Spiridonov and Zhedanov in [17], where the equation is called as just "the discrete-time Toda lattice". On the other hand, equation (2) was proposed by Hirota[5], and called as "the random-time Toda equation". However, it appears that those names are not appropriate for equations (1) and (2), since the former name usually refers the case where λ_t and δ_t are constants, and the latter is somewhat misleading. In this article, we call equations (1) and (2) as "the non-autonomous discrete-time Toda lattice equation".

The non-autonomous discrete-time Toda lattice equation is written in the Lax form

$$L_{t+1}R_{t+1} + \lambda_{t+1} = R_t L_t + \lambda_t, \tag{5}$$

where L_t and R_t are difference operators defined by

$$L_t = A_n^t + e^{-\partial_n}, \quad R_t = B_{n+1}^t e^{\partial_n} + 1, \tag{6}$$

respectively. The Lax equation (5) is the compatibility condition of the spectral problem equation

$$\begin{aligned} \Psi_n^{t+1} &= R_t \Psi_n^t = B_{n+1}^t \Psi_{n+1}^t + \Psi_n^t, \\ (x - \lambda_t)\Psi_n^t &= L_t \Psi_n^{t+1} = A_n^t \Psi_n^{t+1} + \Psi_{n-1}^{t+1}, \end{aligned} \tag{7}$$

where x is a spectral parameter and Ψ_n^t a wave function.

An important feature of soliton equations, including the Toda lattice and the discrete-time Toda equations is that they admit wide class of exact solutions, such as soliton solutions. Moreover, these solutions are expressed by determinants or Pfaffians[6], which are regarded as characteristic property of integrable systems according to the Sato theory[11]. It is known that the discrete-time Toda equation (when λ_t is a constant) admits two kinds of determinant solutions. One is the Hankel type determinant solution, in which the lattice site n appears as the determinant size[3, 7]. Another one is the Casorati determinant solution which describes soliton type solutions[4]. In this solution, the determinant size corresponds to the number of solitons. The Hankel type determinant solution for the non-autonomous discrete-time Toda lattice equation on the semi-infinite lattice was constructed in [12, 13]. The purpose of this article is to present explicit N-soliton solutions for the non-autonomous discrete-time Toda lattice equation in the form of Casorati determinant.

2 Soliton solution for the non-autonomous discrete-time Toda lattice equation

For any $N \in \mathbb{Z}_{>0}$, we first define $N \times N$ Casorati determinants τ_n^t and σ_n^t as

$$\tau_n^t = \begin{vmatrix} \varphi_1^t(n) & \varphi_1^t(n+1) & \cdots & \varphi_1^t(n+N-1) \\ \varphi_2^t(n) & \varphi_2^t(n+1) & \cdots & \varphi_2^t(n+N-1) \\ \vdots & \vdots & & \vdots \\ \varphi_N^t(n) & \varphi_N^t(n+1) & \cdots & \varphi_N^t(n+N-1) \end{vmatrix}, \tag{8}$$

$$\sigma_n^t = \begin{vmatrix} \psi_1^t(n) & \psi_1^t(n+1) & \cdots & \psi_1^t(n+N-1) \\ \psi_2^t(n) & \psi_2^t(n+1) & \cdots & \psi_2^t(n+N-1) \\ \vdots & \vdots & & \vdots \\ \psi_N^t(n) & \psi_N^t(n+1) & \cdots & \psi_N^t(n+N-1) \end{vmatrix}, \tag{9}$$

where the entries $\varphi_i^t(n)$ and $\psi_i^t(n)$ $(i = 1, \ldots, N)$ satisfy linear relations

$$\varphi_i^{t+1}(n) = \varphi_i^t(n) - \mu_t \varphi_i^t(n+1), \tag{10}$$

$$\psi_i^t(n) = \varphi_i^{t-1}(n) - \mu_t \varphi_i^{t-1}(n+1), \tag{11}$$

$$P_i^t \varphi_i^{t-1}(n) = \psi_i^t(n) - \mu_t \psi_i^t(n-1), \tag{12}$$

with μ_t being an arbitrary function of t, and P_i^t given by

$$P_i^t = (1 - p_i \mu_t)(1 - p_i^{-1} \mu_t), \quad i = 1, \ldots, N. \tag{13}$$

For $N = 0$, we put $\tau_n^t = \sigma_n^t = 1$. Then the main result of this article is given as follows:

Theorem 1 *For τ_n^t defined above, the functions*

$$A_n^t = -\mu_t^{-1}\frac{\tau_n^t\tau_{n+1}^{t+1}}{\tau_n^{t+1}\tau_{n+1}^t}, \quad B_n^t = -\mu_t\frac{\tau_{n-1}^{t+1}\tau_{n+1}^t}{\tau_n^t\tau_n^{t+1}}, \quad \lambda_t = \mu_t + \mu_t^{-1}. \tag{14}$$

satisfy the non-autonomous discrete-time Toda lattice equation (1).

As was pointed out in [12, 13], the auxiliary τ function σ_n^t plays an essential role although it does not appear in the final result.

Proposition 2 *τ_n^t and σ_n^t satisfy the following bilinear difference equations:*

$$\tau_n^{t-1}\tau_n^{t+1} - \tau_n^t\sigma_n^t = \mu_{t-1}\mu_t(\tau_{n-1}^{t+1}\tau_{n+1}^{t-1} - \tau_n^t\sigma_n^t), \tag{15}$$

$$\mu_t\sigma_n^t\tau_{n+1}^t - \mu_{t-1}\tau_n^t\sigma_{n+1}^t = (\mu_t - \mu_{t-1})\tau_n^{t+1}\tau_{n+1}^{t-1}. \tag{16}$$

Theorem 1 is a direct consequence of Proposition 2. Actually, multiplying $1-(\mu_t\mu_{t-1})^{-1}$ to equation (16), we have

$$(\mu_t - \mu_{t-1}^{-1})\sigma_n^t\tau_{n+1}^t - (\mu_{t-1} - \mu_t^{-1})\tau_n^t\sigma_{n+1}^t = (\lambda_t - \lambda_{t-1})\tau_n^{t+1}\tau_{n+1}^{t-1}. \tag{17}$$

Multiplying equation (17) by $\tau_n^t\tau_{n+1}^t$ and using equation (15), we have

$$\begin{aligned} & (\tau_{n+1}^t)^2(\mu_t\tau_{n-1}^{t+1}\tau_{n+1}^{t-1} - \mu_{t-1}^{-1}\tau_n^{t-1}\tau_n^{t+1}) - (\tau_n^t)^2(\mu_{t-1}\tau_{n-1}^{t+1}\tau_{n+1}^{t-1} - \mu_t^{-1}\tau_n^{t-1}\tau_n^{t+1}) \\ = \; & (\lambda_t - \lambda_{t-1})\tau_n^t\tau_n^{t+1}\tau_{n+1}^{t-1}\tau_{n+1}^t. \end{aligned} \tag{18}$$

Dividing equation (18) by $\tau_n^t\tau_n^{t+1}\tau_{n+1}^{t-1}\tau_{n+1}^t$, we obtain the first equation of equation (1). The second equation is an identity under the variable transformation (14).

Remark 3 (1) *If we choose the functions $\varphi_i^t(n)$ and $\psi_i^t(n)$ as exponential type functions*

$$\varphi_i^t(n) = \alpha_i p_i^n\prod_{j=t_0}^{t-1}(1-p_i\mu_j) + \beta_i p_i^{-n}\prod_{j=t_0}^{t-1}(1-p_i^{-1}\mu_j), \tag{19}$$

$$\psi_i^t(n) = \alpha_i p_i^n(1-p_i\mu_t)\prod_{j=t_0}^{t-2}(1-p_i\mu_j) + \beta_i p_i^{-n}(1-p_i^{-1}\mu_t)\prod_{j=t_0}^{t-2}(1-p_i^{-1}\mu_j), \tag{20}$$

respectively, where α_i, β_i, p_i $(i=1,\ldots,N)$ are parameters, we have the N-soliton solution. As is shown in [6, 14], the τ fucntions for soliton solutions are expressed as Casorati determinants whose entries are given by exponential type functions.

(2) *In the case where μ_t is a constant, the bilinear equations (15) and (16) reduce to*

$$\tau_n^{t-1}\tau_n^{t+1} - (\tau_n^t)^2 = \mu^2[\tau_{n-1}^{t+1}\tau_{n+1}^{t-1} - (\tau_n^t)^2], \tag{21}$$

which is the bilinear equation of the discrete-time Toda equation[2]. Indeed, the N-soliton solution also reduces to that for discrete-time Toda equation.

(3) *The functions $\varphi_i^t(n)$ $(i=1,\ldots,N)$ satisfy the spectral problem equation*

$$\begin{aligned} & \varphi_i^{t+1}(n) = -\mu_t\varphi_i^t(n+1) + \varphi_i^t(n), \\ & (x_i - \lambda_t)\varphi_i^t(n) = -\mu_t^{-1}\varphi_i^{t+1}(n) + \varphi_i^{t+1}(n-1), \quad x_i = p_i + p_i^{-1}. \end{aligned} \tag{22}$$

Equation (22) is the spectral problem equation (7) with $A_n^t = -\mu_t^{-1}$, $B_n^t = -\mu_t$ and $\lambda_t = \mu_t + \mu_t^{-1}$, which is the simplest solution for the non-autonomous discrete-time Toda lattice equation (1).

3 Proof of Proposition 2

In this section we prove Proposition 2 by using the technique developed in [14, 15]. The bilinear equations (15) and (16) reduce to the Plücker relations, which are quadratic identities among the determinants whose columns are properly shifted. Therefore, we first prepare such difference formulae that express shifted determinants in terms of τ or σ. For simplicity, we introduce notation

$$\tau_n^t = \left| \begin{array}{cccc} 0_t & 1_t & \cdots & (N-1)_t \end{array} \right|, \quad \sigma_n^t = \left| \begin{array}{cccc} \hat{0}_t & \hat{1}_t & \cdots & \widehat{(N-1)}_t \end{array} \right|, \tag{23}$$

where the symbols k_t and $\hat{k}_t$ are column vectors given by

$$k_t = \begin{pmatrix} \varphi_1^t(n+k) \\ \varphi_2^t(n+k) \\ \vdots \\ \varphi_N^t(n+k) \end{pmatrix}, \quad \hat{k}_t = \begin{pmatrix} \psi_1^t(n+k) \\ \psi_2^t(n+k) \\ \vdots \\ \psi_N^t(n+k) \end{pmatrix}, \tag{24}$$

respectively.

Lemma 4 *The following formulae hold:*

$$\tau_n^t = \left| \begin{array}{cccc} 0_t & 1_t & \cdots & (N-2)_t \quad (N-1)_t \end{array} \right|, \tag{25}$$

$$\tau_n^{t-1} = \left| \begin{array}{cccc} 0_t & 1_t & \cdots & (N-2)_t \quad (N-1)_{t-1} \end{array} \right|, \tag{26}$$

$$\mu_{t-1}\tau_n^{t-1} = \left| \begin{array}{cccc} 0_t & 1_t & \cdots & (N-2)_t \quad (N-2)_{t-1} \end{array} \right|, \tag{27}$$

$$(\prod_{i=1}^{N} P_i^t)^{-1}\tau_n^{t+1} = \left| \begin{array}{cccc} \tilde{0}_{t+1} & 1_t & \cdots & (N-2)_t \quad (N-1)_t \end{array} \right|, \tag{28}$$

$$(\prod_{i=1}^{N} P_i^t)^{-1}\mu_t\tau_n^{t+1} = \left| \begin{array}{cccc} \tilde{1}_{t+1} & 1_t & \cdots & (N-2)_t \quad (N-1)_t \end{array} \right|, \tag{29}$$

$$(1-\mu_{t-1}\mu_t)(\prod_{i=1}^{N} P_i^t)^{-1}\sigma_n^t = \left| \begin{array}{cccc} \tilde{0}_{t+1} & 1_t & \cdots & (N-2)_t \quad (N-1)_{t-1} \end{array} \right|, \tag{30}$$

where the symbol $\tilde{k}_t$ is the column vector given by

$$\tilde{k}_t = \begin{pmatrix} (P_1^t)^{-1}\varphi_1^t(n+k) \\ (P_2^t)^{-1}\varphi_2^t(n+k) \\ \vdots \\ (P_N^t)^{-1}\varphi_N^t(n) \end{pmatrix}. \tag{31}$$

Proof of Lemma 4: We first note that $\varphi_i^t(n)$ and $\psi_i^t(n)$ also satisfy the linear relations

$$\varphi_i^{t+1}(n) = \psi_i^t(n) - \mu_{t-1}\psi_i^t(n+1), \tag{32}$$

$$P_i^t\varphi_i^t(n) = \varphi_i^{t+1}(n) - \mu_t\varphi_i^{t+1}(n-1). \tag{33}$$

which follow from equations (10)-(12). Equation (25) is nothing but the definition. Equation (26) is derived as follows: Subtracting $(j+1)$-th column multiplied by μ_{t-1} from j-th column of τ_n^{t-1} for $j = 0, 1, \ldots, N-1$, and using equation (10), we have

$$\begin{aligned} \tau_n^{t-1} &= \left| \begin{array}{cccc} 0_{t-1} & 1_{t-1} & \cdots & (N-1)_{t-1} \end{array} \right| \\ &= \left| \begin{array}{cccc} 0_{t-1} - \mu_{t-1} \times 1_{t-1} & 1_{t-1} & \cdots & (N-1)_{t-1} \end{array} \right| \\ &= \left| \begin{array}{cccc} 0_t & 1_{t-1} & \cdots & (N-1)_{t-1} \end{array} \right| \\ &= \cdots = \left| \begin{array}{cccc} 0_t & \cdots & (N-2)_t & (N-1)_{t-1} \end{array} \right|, \end{aligned}$$

which is equation (26). Moreover, multiplying μ_{t-1} to the N-th column of right hand side of equation (26) and using equation (10), we have

$$\begin{aligned}\mu_{t-1}\tau_n^{t-1} &= \left| \begin{array}{cccc} 1_t & \cdots & (N-2)_t & \mu_{t-1}\times(N-1)_{t-1} \end{array} \right| \\ &= \left| \begin{array}{cccc} 1_t & \cdots & (N-2)_t & (N-2)_t+\mu_{t-1}\times(N-1)_{t-1} \end{array} \right| \\ &= \left| \begin{array}{cccc} 1_t & \cdots & (N-2)_t & (N-2)_{t-1} \end{array} \right|,\end{aligned}$$

which is nothing but equation (27). Equations (28) and (29) can be proved in similar manner by using equation (33). Equation (30) can be proved as follows: first notice that σ_n^t is rewritten as

$$\sigma_n^t = \left| \begin{array}{cccc} 0_{t+1} & \cdots & (N-1)_{t+1} & \widehat{(N-1)}_t \end{array} \right|, \tag{34}$$

which is shown in similar manner by using equation (32). We also note that $\varphi_i^t(n)$ and $\psi_i^t(n)$ satisfy the relation,

$$(1-\mu_t\mu_{t-1})\psi_i^t(n) = P_i^t\varphi_i^{t-1}(n) + \mu_t\varphi_i^{t+1}(n-1), \tag{35}$$

which can be derived by eliminating $\varphi_i^t(n-1)$ from equation (32) with n being replaced by $n-1$ and equation (12). Then multiplying $(1-\mu_t\mu_{t-1})$ to the N-th column of the right hand side of equation (34) and using equation (35), we obtain

$$\begin{aligned}(1-\mu_t\mu_{t-1})\sigma_n^t &= \left| \begin{array}{cccc} 1_{t+1} & \cdots & (N-2)_{t+1} & (1-\mu_t\mu_{t-1})\times\widehat{(N-1)}_{t-1} \end{array} \right| \\ &= \left| \begin{array}{cccc} \varphi_1^{t+1}(n) & \cdots & \varphi_1^{t+1}(n+N-2) & P_1^t\varphi_1^{t-1}(n+N-1) \\ \varphi_2^{t+1}(n) & \cdots & \varphi_2^{t+1}(n+N-2) & P_2^t\varphi_2^{t-1}(n+N-1) \\ \vdots & \cdots & \vdots & \vdots \\ \varphi_N^{t+1}(n) & \cdots & \varphi_N^{t+1}(n+N-2) & P_N^t\varphi_N^{t-1}(n+N-1) \end{array} \right| \\ &= \cdots = \left| \begin{array}{cccc} \varphi_1^{t+1}(n) & P_1^t\varphi_1^{t-1}(n+1) & \cdots & P_1^t\varphi_1^{t-1}(n+N-1) \\ \varphi_2^{t+1}(n) & P_2^t\varphi_2^{t-1}(n+1) & \cdots & P_2^t\varphi_2^{t-1}(n+N-1) \\ \vdots & \vdots & \cdots & \vdots \\ \varphi_N^{t+1}(n) & P_N^t\varphi_N^{t-1}(n+1) & \cdots & P_N^t\varphi_N^{t-1}(n+N-1) \end{array} \right| \\ &= \prod_{i=1}^N P_i^t \left| \begin{array}{cccc} \tilde{0}_{t+1} & 1_{t-1} & \cdots & (N-1)_{t-1} \end{array} \right|,\end{aligned}$$

which is equation (30). This completes the proof of Lemma 4. □

Now consider the following identity of $2N\times 2N$ determinant:

$$\left| \begin{array}{c|ccc|c|cc} \tilde{0}_{t+1} & 0_t & \cdots & (N-2)_t & \multicolumn{1}{c|}{\multirow{2}{*}{$\emptyset$}} & (N-1)_t & (N-1)_{t-1} \\ \tilde{0}_{t+1} & \multicolumn{3}{c|}{} & & (N-1)_t & (N-1)_{t-1} \end{array} \right| = 0, \tag{36}$$

Applying the Laplace expansion to left hand side of equation (36) and using Lemma 4, we obtain

$$\begin{aligned}0 &= \left| \begin{array}{ccccc} \tilde{0}_{t+1} & 0_t & 1_t & \cdots & (N-2)_t \end{array} \right| \times \left| \begin{array}{ccccc} 1_t & \cdots & (N-2)_t & (N-1)_t & (N-1)_{t-1} \end{array} \right| \\ &\quad + \left| \begin{array}{ccccc} 0_t & 1_t & \cdots & (N-2)_t & (N-1)_t \end{array} \right| \times \left| \begin{array}{ccccc} \tilde{0}_{t+1} & 1_t & \cdots & (N-2)_t & (N-1)_{t-1} \end{array} \right| \\ &\quad - \left| \begin{array}{ccccc} 0_t & 1_t & \cdots & (N-2)_t & (N-1)_{t-1} \end{array} \right| \times \left| \begin{array}{ccccc} \tilde{0}_{t+1} & 1_t & \cdots & (N-2)_t & (N-1)_t \end{array} \right| \\ &= \mu_t(\prod_i^N P_i^t)\tau_{n-1}^{t+1}\times\mu_t\tau_{n+1}^{t-1} + \tau_n^t\times(1-\mu_t\mu_{t-1})(\prod_i^N P_i^t)\sigma_n^t - \tau_n^{t-1}\times(\prod_i^N P_i^t)\tau_n^{t+1},\end{aligned}$$

which is the bilinear equation (15).

The bilinear equation (16) can be proved by the similar technique. We prepare the following difference formulae:

Lemma 5 *The following formulae hold.*

$$\tau_n^t = \left| \begin{array}{ccccc} 0_{t+1} & 1_{t+1} & \cdots & (N-2)_{t+1} & (N-1)_t \end{array} \right|, \tag{37}$$

$$\mu_t \tau_{n+1}^t = \left| \begin{array}{ccccc} 1_{t+1} & 2_{t+1} & \cdots & (N-1)_{t+1} & (N-1)_t \end{array} \right|, \tag{38}$$

$$\sigma_n^t = \left| \begin{array}{ccccc} 0_{t+1} & 1_{t+1} & \cdots & (N-2)_{t+1} & \widehat{(N-1)}_t \end{array} \right|, \tag{39}$$

$$\mu_{t-1} \sigma_{n+1}^t = \left| \begin{array}{cccc} 1_{t+1} & \cdots & (N-2)_{t+1} & (N-1)_{t+1} & \widehat{(N-1)}_t \end{array} \right|, \tag{40}$$

$$(\mu_{t-1} - \mu_t) \tau_{n+1}^{t-1} = \left| \begin{array}{cccc} 1_{t+1} & \cdots & (N-2)_{t+1} & (N-1)_t & \widehat{(N-1)}_t \end{array} \right|. \tag{41}$$

Proof. of Lemma 5: Equations (37) and (38) are equivalent to equations (26) and (27), respectively. Equation (39) is the same as equation (34). Equation (40) can be derived by using equation (32) after multiplying μ_{t-1} to the N-th column of the right hand side of equation (37). In order to prove equation (41), we note the following relation between $\varphi_i^t(n)$ and $\psi_i^t(n)$,

$$(\mu_{t-1} - \mu_t) \varphi_i^{t-1}(n) = \psi_i^t(n-1) - \varphi_i^t(n-1), \tag{42}$$

which can be obtained by eliminating $\varphi_i^{t-1}(n)$ from equation (10) with t being replaced by $t-1$ and equation (11). Multiplying $\mu_{t-1} - \mu_t$ on the N-th column of τ_{n+1}^{t-1} and using equation (42), we obtain equation (41). This completes the proof of Lemma 5. □

The bilinear equation (16) is derived by applying the Laplace expansion to left hand side of the following identity

$$\left| \begin{array}{ccc|ccc} 0_{t+1} & \cdots & (N-2)_{t+1} & & \emptyset & & (N-1)_{t+1} & (N-1)_t & \widehat{(N-1)}_t \\ \hline & \emptyset & & 1_{t+1} & \cdots & (N-2)_{t+1} & (N-1)_{t+1} & (N-1)_t & \widehat{(N-1)}_t \end{array} \right| = 0,$$

and using Lemma 5. This completes the proof of Proposition 2 and thus Theorem 1.

4 Concluding remarks

In this article we have presented the N-soliton solution for the non-autonomous discrete-time Toda lattice equation (1), which can be regarded as a generalization of the discrete-time Toda equation such that the lattice interval with respect to time is an arbitrary function in time.

Discrete soliton equations commonly arise as Bäcklund or Darboux type transformations for contiunous soliton equations. In this context, a number of iterations of a Bäcklund transformation can be regarded as the discrete independent variable. The Bäcklund transformation admits one parameter, playing a role of the lattice interval, which can be arbitrary function in correspoiding independent variable. In this sense, discrete soliton equations can be naturally extended to be non-autonomous (see, for example, [1, 16]).

As for solutions, however, it seems that only autonomous cases (the lattice intervals are constants) have been discussed in most of studies. It was recognized in [8, 9] that the discrete two-dimensional Toda lattice equation (or equivalently, the discrete KP equation) admits a non-autonomous generalization. It is known that various discrete soliton equations are derived from the discrete KP equation and its Bäcklund transformations. Therefore it is expected that solutions of non-autonomous discrete soliton equations are discussed from this point of view. For example, the solutions of non-autonomous discrete-time relativistic Toda equation have been constructed in this manner in [10].

However, direct reduction process from the non-autonomous discrete KP equation might not be sufficient. As we have shown in this article, in the case of equation (2), clever introduction of auxiliary τ function (σ_n^t in this article) is critical, which does not appear in the autonomous or continous cases. Careful investigation of this machinery may lead to various generalizations of discrete soliton equations and their solutions. This problem will be discussed in forthcoming articles.

Acknowledgement

The authors would like to thank Dr. Satoshi Tsujimoto and Dr. Ken-ichi Maruno for valuable discussions. One of the author (K.K) thanks Prof. Nalini Joshi for hospitality during his stay in University of Sydney. This work was partly supported by the Grant-in-Aid for JSPS Fellows, The Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

[1] A. I. Bobenko and R. Seiler (eds.), *Discrete integrable geometry and physics*, Oxford Lecture Series in Mathematics and its Applications **16**, Oxford University Press, New York, 1999.

[2] R. Hirota, *Nonlinear partial difference equations. II. Discrete-time Toda equation*, J. Phys. Soc. Japan **43** (1977), 2074–2078.

[3] R. Hirota, *Discrete two-dimensional Toda molecule equation*, J. Phys. Soc. Japan **56** (1987), 4285–4288.

[4] R. Hirota, M. Ito and F. Kako, *Two-dimensional Toda lattice equations*, Prog. Theor. Phys. Suppl. **94** (1988), 42–58.

[5] R. Hirota, *Conserved quantities of "random-time Toda equation"*, J. Phys. Soc. Japan **66** (1997), 283–284.

[6] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge Tracts in Mathematics **155**, Cambridge University Press, New York, 2004.

[7] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *Determinant formulas for the Toda and discrete Toda equations*, Funkcial. Ekvac. **44** (2001), 291–307.

[8] K. Kajiwara and J. Satsuma, *q-difference version of the two-dimensional Toda lattice equation* J. Phys. Soc. Japan **60** (1991), 3986–3989.

[9] K. Kajiwara, Y. Ohta and J. Satsuma, *q-discrete Toda molecule equation*, Phys. Lett. A **180** (1993), 249–256.

[10] K. Maruno, K. Kajiwara and M. Oikawa, *A note on integrable systems related to discrete time Toda lattice*, SIDE III—symmetries and integrability of difference equations (Sabaudia, 1998), CRM Proc. Lecture Notes **25**, Amer. Math. Soc. Providence, 2000, 303–314.

[11] T. Miwa, M. Jimbo and E. Date, *Solitons: Differential equations, symmetries and infinite dimensional algebras*, Cambridge Tracts in Mathematics **135**, Cambridge University Press, New York, 1999.

[12] A. Mukaihira and S. Tsujimoto, *Determinant structure of R_I type discrete integrable system*, J. Phys. A: Math. Gen. **37** (2004), 4557–4565.

[13] A. Mukaihira and S. Tsujimoto, *Determinant structure of spectral transformation chain associated with biorthogonal rational functions*, submitted to J. Phys. A: Math. Gen.

[14] Y. Ohta, R. Hirota, S. Tsujimoto and T. Imai, *Casorati and discrete Gram type determinant representations of solutions to the discrete KP hierarchy*, J. Phys. Soc. Japan **62** (1993), 1872–1886.

[15] Y. Ohta, K. Kajiwara, J. Matsukidaira and J. Satsuma, *Casorati determinant solution for the relativistic Toda lattice equations*, J. Math. Phys. **34** (1993), 5190–5204.

[16] W. K. Schief, *Isothermic surfaces in spaces of arbitrary dimension: integrability, discretization, and Bäcklund transformations—a discrete Calapso equation*, Stud. Appl. Math. **106** (2001), 85–137.

[17] V. Spiridonov and A. Zhedanov, *Discrete Darboux transformations, the discrete-time Toda lattice, and the Askey-Wilson polynomials*, Methods. Appl. Anal. **2** (1995), 369–398.

[18] L. Vinet and A. Zhedanov, *An integrable chain and bi-orthogonal polynomials*, Lett. Math. Phys. **46** (1998), 233–245.